

Changing Mixed Numbers to Improper Fractions

Step one: Multiply the whole number part of the mixed number and the denominator part of the fraction.

Step two: Take the product from step one and add it to the numerator part of the fraction.

Step three: Take the result from step two and write it over the original denominator.

$$\text{Example: } 3 \frac{1}{2} = \frac{3 \cdot 2 + 1}{2} = \frac{7}{2}$$

1. $4 \frac{1}{3} =$	2. $2 \frac{5}{6} =$
3. $3 \frac{4}{7} =$	4. $9 \frac{1}{2} =$
5. $7 \frac{3}{5} =$	6. $4 \frac{3}{4} =$
7. $1 \frac{1}{4} =$	8. $6 \frac{6}{7} =$
9. $5 \frac{7}{9} =$	10. $8 \frac{7}{10} =$

Changing Mixed Numbers to Improper Fractions

Step one: Multiply the whole number part of the mixed number and the denominator part of the fraction.

Step two: Take the product from step one and add it to the numerator part of the fraction.

Step three: Take the result from step two and write it over the original denominator.

$$\text{Example: } 3 \frac{1}{2} = \frac{3 \cdot 2 + 1}{2} = \frac{7}{2}$$

1. $4 \frac{1}{3} = \frac{13}{3}$	2. $2 \frac{5}{6} = \frac{17}{6}$
3. $3 \frac{4}{7} = \frac{25}{7}$	4. $9 \frac{1}{2} = \frac{19}{2}$
5. $7 \frac{3}{5} = \frac{38}{5}$	6. $4 \frac{3}{4} = \frac{19}{4}$
7. $1 \frac{1}{4} = \frac{5}{4}$	8. $6 \frac{6}{7} = \frac{48}{7}$
9. $5 \frac{7}{9} = \frac{52}{9}$	10. $8 \frac{7}{10} = \frac{87}{10}$